

# COVID-19 Vaccine

Monitoring post-vaccination signs and symptoms in LTCF residents

Administered  
in two doses...  
21 (Pfizer) or  
28 (Moderna)  
days apart



95%  
Effective

*Residents and healthcare personnel (HCP) should continue to follow all current CDC infection prevention and control recommendations to protect themselves and others.*

COMMONLY  
REPORTED VACCINE  
**SIDE  
EFFECTS**



FATIGUE



HEADACHE



FEVER



CHILLS



MUSCLE  
ACHE



JOINT  
PAIN

MORE COMMONLY REPORTED  
AFTER 2ND DOSE

Isolate  
and use  
full PPE

Symptoms  
resolve  
w/in 2  
days

DISCONTINUE  
PRECAUTIONS

Symptoms  
persist >2  
days

Isolate  
and use  
full PPE\*


Test for  
SARS-  
CoV-2

Positive  
PCR test

MAINTAIN  
TRANSMISSION  
BASED  
PRECAUTIONS  
AND COHORT

*\* Do NOT cohort with  
SARS-CoV-2 infected  
residents*

SIGNS AND  
SYMPTOMS  
**UNLIKELY  
TO BE VACCINE  
SIDE EFFECTS**



COUGH



SORE THROAT



SHORTNESS  
OF BREATH



RUNNY NOSE



LOSS OF  
TASTE OR  
SMELL

*The vaccine cannot make you sick with COVID-19. The vaccine will not cause you to test positive on viral tests (rapid antigen or PCR).*