

Sepsis Worksheet

Complete for every sepsis patient.

<u>Infection</u>	<u>SIRS</u>	<u>Organ Dysfunction</u>
<input type="checkbox"/> Diff Dx. <input type="checkbox"/> Documented or suspected infection <input type="checkbox"/> ABX ordered for infection	<p style="text-align: center;"><i>Min. required: 2</i></p> <input type="checkbox"/> HR > 90 <input type="checkbox"/> Temp <36.0 or >38.3 C (<96.8 or >100.9 F) <input type="checkbox"/> RR > 20 <input type="checkbox"/> WBC's <4.0 or >12 <u>OR</u> Bands >10	<input type="checkbox"/> Bili >2 <input type="checkbox"/> Lactic Acid >2 <input type="checkbox"/> Cr >2 <input type="checkbox"/> SBP <90 <i>OR</i> MAP <65 <input type="checkbox"/> ACUTE resp. failure requiring vent or BiPap <input type="checkbox"/> Plt <100 <input type="checkbox"/> INR >1.5 <input type="checkbox"/> PTT >60

TIME ZERO: Time that the final sepsis criteria is met within a 6 hr. window

<p style="text-align: center;"><u>Lactic Acid (LA):</u> Drawn within 3 hrs. of time zero</p> <p style="text-align: center;">RN MUST TREND LACTIC ACID #1 and #2 RESULTS. See key below:</p> <p>Initial Lactic Acid Result _____</p> <p><u>Repeat</u> Lactic Acid Result _____ (if ordered / indicated)</p> <p>Key: If LA #2 result is GREATER than LA #1 result center will order additional timed lactic acid q3 hrs x 2 (protocol / no cosign required)</p> <p>Third Lactic Acid Result _____ (if ordered / indicated)</p>	<p>Date/time & Emp #</p>
<p style="text-align: center;"><u>Blood Cx:</u> Drawn within 3 hrs. of time zero</p> <p style="text-align: center;">Key: DRAW PERIPHERALLY EVEN IF ALREADY ON ABX</p>	
<p style="text-align: center;"><u>IV Antibiotics:</u> Admin. within 3 hrs. of time zero (or up to 6 hrs before) (may admin IM/IO if limited IV access is documented)</p> <p style="text-align: center;">Keys: 1. ALWAYS admin. Broad spectrum ABX 1st when multiple. ABX ordered</p>	
<p style="text-align: center;"><u>Sepsis IV Bolus (LR or NS):</u></p> <p style="text-align: center;">Required for:</p> <p style="text-align: center;">Lactic Acid: 4 or more</p> <p style="text-align: center;">OR</p> <p style="text-align: center;">MAP <65 <i>OR</i> SBP <90 x 2 readings in a 3 hr window</p> <p style="text-align: center;">Key: All pts receiving a sepsis bolus <u>MUST</u> have VS q30 minutes x 2 hrs. <u>AFTER</u> initiating the final bolus volume.</p>	
<p><u>Pressors:</u> Required for new onset or persistent hypotension present in the 1 hour window after the crystalloid sepsis bolus completion.</p>	